

## WORLD INTELLECTUAL PROPERTY ORGANIZA N International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> :	Δ1	(11) International Publication Number:	WO 98/08991
C22B 23/00		(43) International Publication Date:	5 March 1998 (05.03.98)

(21) International Application Number: PCT/US97/15109

(22) International Filing Date: 29 August 1997 (29.08.97)

(30) Priority Data: 60/024,928 30 August 1996 (30.08.96) US 60/030,462 6 November 1996 (06.11.96) US

(71)(72) Applicants and Inventors: CHANEY, Rufus, L. [US/US]; United States Department of Agriculture, Beltsville, MD 20705 (US). ANGLE, Jay, Scott [US/US]; 10241 Bristol Channel, Ellicot City, MA 21042 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): LI, Yin-Ming [CN/US]; 12019 Coldstream Drive, Potomac, MD 20854 (US).

(74) Agents: KELBER, Steven, B. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, P.C., Crystal Square Five, 4th floor, 1755 Jefferson Davis Highway, Arlington, VA 22202 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TT, TM, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

## **Published**

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: METHOD FOR PHYTOMINING OF NICKEL, COBALT AND OTHER METALS FROM SOIL

## (57) Abstract

The recovery of nickel, cobalt and other metals by phytomining is described. Plants of the Alyssum genus are grown in nickel rich soil. The uptake of nickel is enhanced by maintaining specific soil conditions, including a concentration of calcium between (but not including) 0.128 mM and 5.0 mM and an acidic pH. Ni uptake may be further enhanced by maintaining a ratio of exchangeable Ca/Mg of 0.16-0.40. Uptake may be further enhanced by addition of chelating agents and ammonium based fertilizers.

